

Geometry formulas (given on the exam if needed)

Rectangle Area = base \times height

Triangle Area = $\frac{1}{2}$ base \times height.

Perimeter of a polygon (triangle, rectangle, etc.) = sum of side lengths

Circumference of a circle = $2\pi r$

Area of a circle = πr^2

Surface area of a rectangular box = sum of all sides' areas

Volume of a rectangular box = length \times width \times height

Surface area of a cylinder = $2\pi r^2 + 2\pi r h$

Volume of a cylinder = $\pi r^2 h$

Pythagorean theorem: $a^2 + b^2 = c^2$

Useful formulas

Linear functions: $y = mx + b$ (slope-intercept form), $y - y_1 = m(x - x_1)$ (point-slope form)

Profit, Revenue, and Cost: Profit = Revenue - Cost

Quadratic Formula: $ax^2 + bx + c = 0 \rightsquigarrow x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$

Five derivative rules for operations on functions.

Constant Multiple Rule: $\frac{d}{dx}(cf(x)) = cf'(x)$

Sum and Difference Rule: $\frac{d}{dx}(f(x) \pm g(x)) = f'(x) \pm g'(x)$

Product Rule: $\frac{d}{dx}(f(x) \cdot g(x)) = f'(x)g(x) + f(x)g'(x)$

Quotient Rule: $\frac{d}{dx}\left(\frac{f(x)}{g(x)}\right) = \frac{f'(x)g(x) - f(x)g'(x)}{(g(x))^2}$

Chain Rule: $\frac{d}{dx}(f(g(x))) = f'(g(x)) \cdot g'(x)$

Ten derivative shortcuts for functions

Derivative of a Constant: $\frac{d}{dx}(c) = 0$, where c is a constant.

The Power Rule: $\frac{d}{dx}(x^n) = nx^{n-1}$

Exponential Functions: $\frac{d}{dx}(a^x) = a^x \cdot \ln(a)$

Special Case: $\frac{d}{dx}(e^x) = e^x$

Three Trigonometric Rules:

$$\frac{d}{dx}(\sin(x)) = \cos(x)$$

$$\frac{d}{dx}(\cos(x)) = -\sin(x)$$

$$\frac{d}{dx}(\tan(x)) = \sec^2(x) = \frac{1}{\cos^2(x)}$$

Three Inverse Function Rules:

$$\frac{d}{dx}(\ln(x)) = \frac{1}{x}$$

$$\frac{d}{dx}(\arctan(x)) = \frac{1}{1+x^2}$$

$$\frac{d}{dx}(\arcsin(x)) = \frac{1}{\sqrt{1-x^2}}$$

General antiderivative rules

If k is a constant $\int k dx = kx + C$

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C, \text{ when } n \neq -1$$

$$\int a^x dx = \frac{a^x}{\ln(a)} + C$$

$$\int e^x dx = e^x + C$$

$$\int \cos(x) dx = \sin(x) + C$$

$$\int \sin(x) dx = -\cos(x) + C$$

$$\int \sec^2(x) dx = \tan(x) + C$$

$$\int \frac{1}{x} dx = \ln(|x|) + C$$

$$\int \frac{1}{1+x^2} dx = \arctan(x) + C$$

$$\int \frac{1}{\sqrt{1-x^2}} dx = \arcsin(x) + C$$